

Product Data Sheet

PRORSD1 siRNA (Mouse)

Catalog #	Source	Reactivity	Applic	cations	
CRM7434	Synthetic	Μ	RNAi		
Description	siRNA	to inhibit PRORSD1	expression using RNA in	terference	
Specificity	PROR	SD1 siRNA (Mouse) is	a target-specific 19-23	nt siRNA oligo duplexes designed	
	to kno	ock down gene expre	ssion.		
Form	Lyoph	nilized powder			
Gene Symbol	PROR	PRORSD1			
Alternative N	ames PRDX	PRDXDD1; Prolyl-tRNA synthetase associated domain-containing protein 1; PrdX			
	deacy	lase domain-containi	ng protein 1		
Entrez Gene	67939	9 (Mouse)			
SwissProt Q9D820 (Mouse)					
Purity	> 97%				
Quality Contr	ol Oligoi	Oligonucleotide synthesis is monitored base by base through trityl analysis to en			
	appro	opriate coupling efficie	ency. The oligo is subse	quently purified by affinity-solid	
	phase	e extraction. The anne	ealed RNA duplex is furt	her analyzed by mass	
	spect	rometry to verify the	exact composition of th	ne duplex. Each lot is compared to	
	the p	revious lot by mass sp	ectrometry to ensure r	naximum lot-to-lot consistency.	
Components We offers pre-designed sets of 3 different target-specific siRNA oligo duple			pecific siRNA oligo duplexes of		
	mous	e PRORSD1 gene. Eac	h vial contains 5 nmol o	of lyophilized siRNA. The duplexes	
	can b	e transfected individu	ally or pooled together	to achieve knockdown of the	
	target	t gene, which is most	commonly assessed by	qPCR or western blot.	
	Com	ponent	15 nmo	l 30 nmol	
	PRO	RSD1 siRNA (Mouse)	- A 5 nmol	x 1 5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

PRORSD1 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
PRORSD1 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 μl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 μl
		10 nM	1 μl	5 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com