

OOEP siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications |
|--------------------------|---|------------|--------------|
| CRM7450 | Synthetic | M | RNAi |
| Description | siRNA to inhibit OOEP expression using RNA interference | | |
| Specificity | OOEP siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to knock down gene expression. | | |
| Form | Lyophilized powder | | |
| Gene Symbol | OOEP | | |
| Alternative Names | OEP19; SDDR; Oocyte-expressed protein homolog; Factor located in oocytes permitting embryonic development; Floped; Oocyte- and embryo-specific protein 19; mOEP19; STAT3 downstream gene and differentiation regulator | | |
| Entrez Gene | 67968 (Mouse) | | |
| SwissProt | Q9CWE6 (Mouse) | | |
| Purity | > 97% | | |
| Quality Control | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency. | | |
| Components | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of mouse OOEP gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can be transfected individually or pooled together to achieve knockdown of the target gene, which is most commonly assessed by qPCR or western blot. | | |

| Component | 15 nmol | 30 nmol |
|-----------|---------|---------|
|-----------|---------|---------|

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB
www.cohesionbio.com

ORDER
order@cohesionbio.com

SUPPORT
techsupport@cohesionbio.com

CUSTOM
custom@cohesionbio.com

Product Data Sheet

| | | |
|------------------------|--------------|--------------|
| OOEP siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 |
| OOEP siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
| OOEP siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume of medium | Final concentration of siRNA | siRNA (20 μ M) | Lipofectamin 2000 |
|---------|------------------------|------------------------------|--------------------|-------------------|
| 96-well | 100 μ l | 100 nM | 0.5 μ l | 0.25 μ l |
| | | 50 nM | 0.25 μ l | 0.25 μ l |
| | | 10 nM | 0.05 μ l | 0.25 μ l |
| 24-well | 500 μ l | 100 nM | 2.5 μ l | 1 μ l |
| | | 50 nM | 1.25 μ l | 1 μ l |
| | | 10 nM | 0.25 μ l | 1 μ l |
| 12-well | 1 ml | 100 nM | 5 μ l | 2 μ l |
| | | 50 nM | 2.5 μ l | 2 μ l |
| | | 10 nM | 0.5 μ l | 2 μ l |
| 6-well | 2 ml | 100 nM | 10 μ l | 5 μ l |
| | | 50 nM | 5 μ l | 5 μ l |
| | | 10 nM | 1 μ l | 5 μ l |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB

www.cohesionbio.com

ORDER

order@cohesionbio.com

SUPPORT

techsupport@cohesionbio.com

CUSTOM

custom@cohesionbio.com

Product Data Sheet

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, CHIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB

www.cohesionbio.com

ORDER

order@cohesionbio.com

SUPPORT

techsupport@cohesionbio.com

CUSTOM

custom@cohesionbio.com