

Product Data Sheet

SLC25A22 siRNA (Mouse)

Catalog #	Source	Reactivity		Applications		
CRM7586	Synthetic	М	I	RNAi		
Description	siRNA	to inhibit SLC25A22	expression using l	RNA interference		
Specificity	SLC25	5A22 siRNA (Mouse) i	s a target-specific	19-23 nt siRNA oligo duplexes designed		
	to kno	ock down gene expre	ssion.			
Form	Lyoph	nilized powder				
Gene Symbol	SLC25	SLC25A22				
Alternative N	ames GC1;	GC1; Mitochondrial glutamate carrier 1; GC-1; Glutamate/H(+) symporter 1; Solute				
	carrie	r family 25 member 2	22			
Entrez Gene	68267	7 (Mouse)				
SwissProt Q9D6M3 (Mouse)						
Purity	> 97%	0				
Quality Control Oligonucleotide synthesis is monitored base by base through trityl ana			by base through trityl analysis to ensure			
	appro	opriate coupling efficie	ency. The oligo is	subsequently purified by affinity-solid		
	phase	e extraction. The anne	ealed RNA duplex	is further analyzed by mass		
	spect	rometry to verify the	exact composition	n of the duplex. Each lot is compared to		
	the p	revious lot by mass sp	pectrometry to en	sure maximum lot-to-lot consistency.		
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	mous	e SLC25A22 gene. Ead	ch vial contains 5	nmol of lyophilized siRNA. The duplexes		
	can b	e transfected individu	ally or pooled tog	gether to achieve knockdown of the		
	target	target gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	1	5 nmol 30 nmol		
	SLC2	5A22 siRNA (Mouse)	- A 5	nmol x 1 5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

SLC25A22 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
SLC25A22 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 µl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
_		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 μΙ
		10 nM	1 μΙ	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com