

# **Product Data Sheet**

## PIK3CB siRNA (Mouse)

| Catalog #     | Source    | Reactivity   | Applications   |  |  |
|---------------|-----------|--|--|--|--|
| CRM9341       | Synthetic | М  | RNAi   |  |  |
| Description   | siRN      | A to inhibit PIK3CB ex   | pression using RNA interference  |  |  |
| Specificity   | PIK3      | CB siRNA (Mouse) is a  | target-specific 19-23 nt siRNA oligo duplexes designed to  |  |  |
|               | knoc      | k down gene expressi   | on.  |  |  |
| Form          | Lyop      | hilized powder   |  |  |  |
| Gene Symbol   | PIK3      | PIK3CB   |  |  |  |
| Alternative N | ames Phos | Phosphatidylinositol 4.5-bisphosphate 3-kinase catalytic subunit beta isoform; |  |  |  |
|               | PI3-k     | kinase subunit beta; P   | I3K-beta; PI3Kbeta; PtdIns-3-kinase subunit beta;  |  |  |
|               | Phos      | phatidylinositol 4.5-b   | isphosphate 3-kinase 110 kDa catalytic subunit beta;   |  |  |
|               | PtdIr     | ns-3-kinase subun  |  |  |  |
| Entrez Gene   | 7476      | 74769 (Mouse)  |  |  |  |
| SwissProt     | Q8B       | ΓΙ9 (Mouse)  |  |  |  |
| Purity        | > 979     | %  |  |  |  |
| Quality Contr | ol Oligo  | onucleotide synthesis  | is monitored base by base through trityl analysis to ensure  |  |  |
|               | appr      | opriate coupling effici  | ency. The oligo is subsequently purified by affinity-solid   |  |  |
|               | phas      | e extraction. The ann  | ealed RNA duplex is further analyzed by mass   |  |  |
|               | spec      | trometry to verify the   | exact composition of the duplex. Each lot is compared to   |  |  |
|               | the p     | previous lot by mass s   | pectrometry to ensure maximum lot-to-lot consistency.  |  |  |
| Components    | We c      | offers pre-designed se   | ts of 3 different target-specific siRNA oligo duplexes of  |  |  |
|               | mou       | se PIK3CB gene. Each   | vial contains 5 nmol of lyophilized siRNA. The duplexes  |  |  |
|               | can b     | be transfected individ   | ually or pooled together to achieve knockdown of the   |  |  |
|               | targe     | et gene, which is most   | commonly assessed by qPCR or western blot.   |  |  |
|               |           |  | - historic sector in the sector of the secto |  |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

| Component                | 15 nmol      | 30 nmol      |
|--------------------------|--------------|--------------|
| PIK3CB siRNA (Mouse) - A | 5 nmol x 1   | 5 nmol x 2   |
| PIK3CB siRNA (Mouse) - B | 5 nmol x 1   | 5 nmol x 2   |
| PIK3CB siRNA (Mouse) - C | 5 nmol x 1   | 5 nmol x 2   |
| Negative Control         | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water               | 1 ml x 1     | 1 ml x 2     |

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate   | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
|         | of medium    | of siRNA            |               | 2000         |
|         |              | 100 nM              | 0.5 μl        | 0.25 μl      |
| 96-well | 100 μl       | 50 nM               | 0.25 μl       | 0.25 μl      |
|         |              | 10 nM               | 0.05 μl       | 0.25 μl      |
|         |              | 100 nM              | 2.5 μl        | 1 μΙ         |
| 24-well | 500 μl       | 50 nM               | 1.25 μl       | 1 μΙ         |
|         |              | 10 nM               | 0.25 μl       | 1 μΙ         |
|         |              | 100 nM              | 5 µl          | 2 μΙ         |
| 12-well | 1 ml         | 50 nM               | 2.5 μl        | 2 μΙ         |
|         |              | 10 nM               | 0.5 μl        | 2 μΙ         |
| 6-well  | 2 ml         | 100 nM              | 10 µl         | 5 µl         |
|         |              | 50 nM               | 5 µl          | 5 μΙ         |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com