

Product Data Sheet

RAI14 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | |
|---------------|------------|--|--|--|--|
| CRM9483 | Synthetic | М | RNAi | | |
| Description | siRNA | A to inhibit RAI14 exp | ression using RNA interference | | |
| Specificity | RAI14 | 4 siRNA (Mouse) is a t | arget-specific 19-23 nt siRNA oligo duplexes designed to | | |
| | knocl | k down gene expressi | on. | | |
| Form | Lyoph | nilized powder | | | |
| Gene Symbol | RAI14 | RAI14 | | | |
| Alternative N | ames KIAA: | KIAA1334; NORPEG; Ankycorbin; Ankyrin repeat and coiled-coil structure-containing | | | |
| | prote | in; Novel retinal pigm | ent epithelial cell protein; Retinoic acid-induced protein | | |
| | 14; p | 125 | | | |
| Entrez Gene | 7564 | 6 (Mouse) | | | |
| SwissProt | Q9EP | 71 (Mouse) | | | |
| Purity | > 97% | 6 | | | |
| Quality Contr | ol Oligo | nucleotide synthesis | s monitored base by base through trityl analysis to ensure | | |
| | appro | opriate coupling effici | ency. The oligo is subsequently purified by affinity-solid | | |
| | phase | e extraction. The anne | aled RNA duplex is further analyzed by mass | | |
| | spect | rometry to verify the | exact composition of the duplex. Each lot is compared to | | |
| | the p | revious lot by mass sp | ectrometry to ensure maximum lot-to-lot consistency. | | |
| Components | We o | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | mous | se RAI14 gene. Each v | al contains 5 nmol of lyophilized siRNA. The duplexes can | | |
| | be tra | be transfected individually or pooled together to achieve knockdown of the target | | | |
| | gene, | gene, which is most commonly assessed by qPCR or western blot. | | | |
| | Com | nponent | 15 nmol 30 nmol | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| RAI14 siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 |
|-------------------------|--------------|--------------|
| RAI14 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
| RAI14 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μΙ |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μl |
| | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 µl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μl |
| | | 10 nM | 1 μl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

Product Data Sheet

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com