

Product Data Sheet

PLEKHJ1 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | |
|---------------|-----------|---|----------------------------------|----------------------------|--|
| CRM9952 | Synthetic | М | RNAi | | |
| Description | siRNA | to inhibit PLEKHJ1 ex | pression using RNA interferenc | e | |
| Specificity | PLEKH | PLEKHJ1 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed | | | |
| | to kno | ock down gene expres | sion. | | |
| Form | Lyoph | ilized powder | | | |
| Gene Symbol | PLEKH | PLEKHJ1 | | | |
| Alternative N | ames GNRP | GNRPX; Pleckstrin homology domain-containing family J member 1; PH | | | |
| | domai | in-containing family J | member 1; Guanine nucleotide | e-releasing protein x | |
| Entrez Gene | 78670 | (Mouse) | | | |
| SwissProt | Q9D24 | Q9D240 (Mouse) | | | |
| Purity | > 97% | > 97% | | | |
| Quality Contr | ol Oligon | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | |
| | appro | priate coupling efficie | ency. The oligo is subsequently | purified by affinity-solid | |
| | phase | extraction. The anne | aled RNA duplex is further anal | yzed by mass | |
| | spectr | ometry to verify the | exact composition of the duple: | x. Each lot is compared to | |
| | the pr | evious lot by mass sp | ectrometry to ensure maximun | n lot-to-lot consistency. | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | mouse | e PLEKHJ1 gene. Each | vial contains 5 nmol of lyophili | zed siRNA. The duplexes | |
| | can be | e transfected individu | ally or pooled together to achie | eve knockdown of the | |
| | target | target gene, which is most commonly assessed by qPCR or western blot. | | | |
| | Com | ponent | 15 nmol | 30 nmol | |
| | PLEK | HJ1 siRNA (Mouse) - / | A 5 nmol x 1 | 5 nmol x 2 | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| PLEKHJ1 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
|---------------------------|--------------|--------------|
| PLEKHJ1 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 μl | 2 µl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 µl |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 µl | 5 μΙ |
| | | 10 nM | 1 μΙ | 5 μΙ |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |