

# **Product Data Sheet**

### **ARL4D siRNA (Mouse)**

Catalog #	Source	Reactivity	Applications			
CRN0106	Synthetic	М	RNAi			
Description	siRNA	to inhibit ARL4D exp	pression using RNA interference			
Specificity	ARL4I	D siRNA (Mouse) is a	target-specific 19-23 nt siRNA oli្	go duplexes designed to		
	knock	down gene expressi	on.			
Form	Lyoph	nilized powder				
Gene Symbol	ARL4I	ARL4D				
Alternative N	ames ARF4I	ARF4L; ARL5; ADP-ribosylation factor-like protein 4D; ADP-ribosylation factor-like				
	prote	in 4L; ADP-ribosylatic	on factor-like protein 5			
Entrez Gene	80982	1 (Mouse)				
SwissProt Q99		Q99PE9 (Mouse)				
Purity > 97%		97%				
Quality Contr	ol Oligoi	Oligonucleotide synthesis is monitored base by base through trityl analysis to e				
	appro	priate coupling effici	ency. The oligo is subsequently p	urified by affinity-solid		
	phase	e extraction. The anne	ealed RNA duplex is further analy	zed by mass		
	spect	rometry to verify the	exact composition of the duplex.	Each lot is compared to		
	the p	revious lot by mass sp	pectrometry to ensure maximum	lot-to-lot consistency.		
Components We offers pre-designed sets of 3 different			ts of 3 different target-specific siR	t target-specific siRNA oligo duplexes of		
	mous	e ARL4D gene. Each v	vial contains 5 nmol of lyophilized	siRNA. The duplexes can		
	be tra	ansfected individually	or pooled together to achieve kn	ockdown of the target		
	gene,	gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
	ARL	4D siRNA (Mouse) - A	5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



# **Product Data Sheet**

ARL4D siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
ARL4D siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 μl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μl
_		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 μΙ	5 μΙ

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com