

# **Product Data Sheet**

## GALNT7 siRNA (Mouse)

| Catalog #                         | Source     | Reactivity  | Applications                          |                           |  |
|-----------------------------------|------------|---|---------------------------------------|---------------------------|--|
| CRN0928                           | Synthetic  | Μ   | RNAi                                  |                           |  |
| Description                       | siRNA      | to inhibit GALNT7 ex  | pression using RNA interference       |                           |  |
| Specificity                       | GALN       | T7 siRNA (Mouse) is   | a target-specific 19-23 nt siRNA o    | ligo duplexes designed to |  |
|                                   | knock      | down gene expressi  | on.                                   |                           |  |
| Form                              | Lyoph      | ilized powder   |                                       |                           |  |
| Gene Symbol                       | GALN       | GALNT7  |                                       |                           |  |
| Alternative N                     | ames N-ace | N-acetylgalactosaminyltransferase 7; Polypeptide GalNAc transferase 7; GalNAc-T7;     |                                       |                           |  |
|                                   | pp-Ga      | pp-GaNTase 7; Protein-UDP acetylgalactosaminyltransferase 7;                          |                                       |                           |  |
|                                   | UDP-0      | GalNAc:polypeptide N  | N-acetylgalactosaminyltransferase     | 2 7                       |  |
| Entrez Gene                       | 10815      | 108150 (Mouse)  |                                       |                           |  |
| SwissProt                         | Q80V/      | Q80VA0 (Mouse)  |                                       |                           |  |
| Purity                            | > 97%      | > 97%   |                                       |                           |  |
| Quality Contr                     | ol Oligor  | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure |                                       |                           |  |
|                                   | appro      | priate coupling effici  | ency. The oligo is subsequently p     | urified by affinity-solid |  |
|                                   | phase      | extraction. The anno  | ealed RNA duplex is further analy     | zed by mass               |  |
|                                   | spectr     | rometry to verify the   | exact composition of the duplex.      | Each lot is compared to   |  |
|                                   | the pr     | revious lot by mass sp  | pectrometry to ensure maximum         | lot-to-lot consistency.   |  |
| Components We offers pre-designed |            |   | ts of 3 different target-specific siR | NA oligo duplexes of      |  |
|                                   | mouse      | e GALNT7 gene. Each   | vial contains 5 nmol of lyophilize    | ed siRNA. The duplexes    |  |
|                                   | can be     | e transfected individu  | ally or pooled together to achiev     | e knockdown of the        |  |
|                                   | target     | gene, which is most   | commonly assessed by qPCR or v        | western blot.             |  |
|                                   | Com        | ponent  | 15 nmol                               | 30 nmol                   |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



# **Product Data Sheet**

| GALNT7 siRNA (Mouse) - A | 5 nmol x 1   | 5 nmol x 2   |
|--------------------------|--------------|--------------|
| GALNT7 siRNA (Mouse) - B | 5 nmol x 1   | 5 nmol x 2   |
| GALNT7 siRNA (Mouse) - C | 5 nmol x 1   | 5 nmol x 2   |
| Negative Control         | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water               | 1 ml x 1     | 1 ml x 2     |

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate   | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
|         | of medium    | of siRNA            |               | 2000         |
|         |              | 100 nM              | 0.5 μl        | 0.25 μl      |
| 96-well | 100 µl       | 50 nM               | 0.25 μl       | 0.25 μl      |
|         |              | 10 nM               | 0.05 μl       | 0.25 μl      |
|         |              | 100 nM              | 2.5 μl        | 1 μΙ         |
| 24-well | 500 μl       | 50 nM               | 1.25 μl       | 1 μΙ         |
|         |              | 10 nM               | 0.25 μl       | 1 μl         |
|         |              | 100 nM              | 5 μl          | 2 μΙ         |
| 12-well | 1 ml         | 50 nM               | 2.5 μl        | 2 μΙ         |
|         |              | 10 nM               | 0.5 μl        | 2 μΙ         |
|         |              | 100 nM              | 10 µl         | 5 µl         |
| 6-well  | 2 ml         | 50 nM               | 5 μl          | 5 μl         |
|         |              | 10 nM               | 1 μl          | 5 μΙ         |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

## **Product Data Sheet**

**Storage/Stability** Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com