

Product Data Sheet

FOXP2 siRNA (Mouse)

Catalog #	Source	Reactivity	Applications		
CRN1176	Synthetic	Μ	RNAi		
Description	siRNA	to inhibit FOXP2 exp	ression using RNA interference		
Specificity	FOXP2	FOXP2 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knock	down gene expressio	on.		
Form	Lyoph	ilized powder			
Gene Symbol	FOXP2	FOXP2			
Alternative Na	ames Forkh	ead box protein P2			
Entrez Gene	11414	2 (Mouse)			
SwissProt	P5846	53 (Mouse)			
Purity >		> 97%			
Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis			rityl analysis to ensure		
	appro	priate coupling efficie	ency. The oligo is subsequently puri	fied by affinity-solid	
	phase	extraction. The anne	aled RNA duplex is further analyzed	d by mass	
	spectr	rometry to verify the	exact composition of the duplex. Ea	ach lot is compared to	
	the pr	evious lot by mass sp	ectrometry to ensure maximum lot	-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	e FOXP2 gene. Each v	ial contains 5 nmol of lyophilized sif	RNA. The duplexes can	
	be tra	nsfected individually	or pooled together to achieve knoc	kdown of the target	
	gene, which is most commonly assessed by qPCR or western blot.			lot.	
	Com	ponent	15 nmol	30 nmol	
	FOXF	2 siRNA (Mouse) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

5 nmol x 1

5 nmol x 2

FOXP2 siRNA (Mouse) - B

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μΙ	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com