

Product Data Sheet

HSD17B11 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | |
|----------------|-----------|---|---|--|--|
| CRN1202 | Synthetic | Μ | RNAi | | |
| Description | siRNA | A to inhibit HSD17B1 | 1 expression using RNA interference | | |
| Specificity | HSD1 | 7B11 siRNA (Mouse) | is a target-specific 19-23 nt siRNA oligo duplexes designed | | |
| | to kn | ock down gene expr | ession. | | |
| Form | Lyopl | nilized powder | | | |
| Gene Symbol | HSD1 | 7B11 | | | |
| Alternative Na | ames DHRS | DHRS8; PAN1B; Estradiol 17-beta-dehydrogenase 11; 17-beta-hydroxysteroid | | | |
| | dehy | drogenase 11; 17-be | ta-HSD 11; 17bHSD11; 17betaHSD11; | | |
| | 17-be | eta-hydroxysteroid d | ehydrogenase XI; 17-beta-HSD XI; 17betaHSDXI; | | |
| | Dehy | drogenase/reductase | e SDR family member 8 | | |
| Entrez Gene | 1146 | 64 (Mouse) | | | |
| SwissProt | Q9EC | Q06 (Mouse) | | | |
| Purity | > 97% | 6 | | | |
| Quality Contro | ol Oligo | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | |
| | appro | opriate coupling effic | iency. The oligo is subsequently purified by affinity-solid | | |
| | phase | e extraction. The anr | ealed RNA duplex is further analyzed by mass | | |
| | spect | rometry to verify the | e exact composition of the duplex. Each lot is compared to | | |
| | the p | revious lot by mass s | pectrometry to ensure maximum lot-to-lot consistency. | | |
| Components | We o | ffers pre-designed se | ets of 3 different target-specific siRNA oligo duplexes of | | |
| | mous | se HSD17B11 gene. E | ach vial contains 5 nmol of lyophilized siRNA. The duplexes | | |
| | can b | e transfected individ | ually or pooled together to achieve knockdown of the | | |
| | targe | t gene, which is mos | t commonly assessed by qPCR or western blot. | | |
| | | | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| Component | 15 nmol | 30 nmol |
|----------------------------|--------------|--------------|
| HSD17B11 siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 |
| HSD17B11 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
| HSD17B11 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μΙ |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 µl | 2 µl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| 6 woll | 2 ml | 100 nM | 10 µl | 5 µl |
| 6-well | Z mi | 50 nM | 5 µl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com