

# **Product Data Sheet**

### CCDC64B siRNA (Mouse)

Catalog #	Source	Reactivity	Applicatio	ons	
CRN1794	Synthetic	Μ	RNAi		
Description	siRNA	to inhibit CCDC64B e	xpression using RNA interf	erence	
Specificity	CCDC	64B siRNA (Mouse) is	a target-specific 19-23 nt s	iRNA oligo duplexes designed	
	to kno	ock down gene expres	sion.		
Form	Lyoph	ilized powder			
Gene Symbol	CCDC	64B			
Alternative N	ames BICDR	BICDR2; Bicaudal D-related protein 2; BICD-related protein 2; BICDR-2; Coiled-coil			
	doma	in-containing protein	64B		
Entrez Gene	21273	33 (Mouse)			
SwissProt	Q8CH	Q8CHW5 (Mouse)			
Purity	> 97%	> 97%			
Quality Contr	ol Oligor	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appro	priate coupling efficie	ency. The oligo is subseque	ntly purified by affinity-solid	
	phase	extraction. The anne	aled RNA duplex is further	analyzed by mass	
	specti	rometry to verify the	exact composition of the d	uplex. Each lot is compared to	
	the pr	evious lot by mass sp	ectrometry to ensure max	imum lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	e CCDC64B gene. Eac	n vial contains 5 nmol of ly	ophilized siRNA. The duplexes	
	can be	e transfected individu	ally or pooled together to	achieve knockdown of the	
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	CCDO	C64B siRNA (Mouse) -	A 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

CCDC64B siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
CCDC64B siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 μl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 μΙ	5 μΙ

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com