

# **Product Data Sheet**

### TIMM8A2 siRNA (Mouse)

Catalog #	Source	Reactivity	Applicat	ions		
CRN2280	Synthetic	М	RNAi			
Description	Description siRNA to inhibit TIMM8A2 expression using RNA interference					
Specificity	TIMM	TIMM8A2 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed				
	to kno	ock down gene expre	ssion.			
Form	Lyoph	nilized powder				
Gene Symbol	TIMM	TIMM8A2				
Alternative N	ames TIMM	TIMM8AB; Putative mitochondrial import inner membrane translocase subunit Tim8				
	A-B					
Entrez Gene	22320	223262 (Mouse)				
SwissProt	Q4FZ	Q4FZG7 (Mouse)				
Purity	> 97%	6				
Quality Contr	Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis to			through trityl analysis to ensure		
	appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli				
	phase	e extraction. The anne	ealed RNA duplex is furthe	er analyzed by mass		
	spect	rometry to verify the	exact composition of the	duplex. Each lot is compared to		
	the p	revious lot by mass sp	ectrometry to ensure ma	ximum lot-to-lot consistency.		
Components We offers pre-designed sets of 3 different target-specific siRNA oligo duplexe			cific siRNA oligo duplexes of			
	mous	e TIMM8A2 gene. Ea	ch vial contains 5 nmol of	lyophilized siRNA. The duplexes		
	can b	e transfected individu	ally or pooled together to	achieve knockdown of the		
	target	t gene, which is most	commonly assessed by q	PCR or western blot.		
	Com	ponent	15 nmol	30 nmol		
	TIMI	M8A2 siRNA (Mouse)	- A 5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

TIMM8A2 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
TIMM8A2 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 µl	5 μΙ

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com