

Product Data Sheet

MARCH2 siRNA (Mouse)

Catalog #	Source	Reactivity	Applications		
CRN2379	Synthetic	М	RNAi		
Description	siRNA	siRNA to inhibit MARCH2 expression using RNA interference			
Specificity	MARC	MARCH2 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed			
	to kno	ock down gene expre	ession.		
Form	Lyoph	ilized powder			
Gene Symbol	MARC	MARCH2			
Alternative N	ames E3 ub	E3 ubiquitin-protein ligase MARCH2; Membrane-associated RING finger protein 2;			
	Meml	brane-associated RIN	IG-CH protein II; MARCH-II		
Entrez Gene	22470	224703 (Mouse)			
SwissProt Q99M02 (Mouse)					
Purity > 97%					
Quality Contr	ol Oligoi	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appro	priate coupling effic	iency. The oligo is subsequently	purified by affinity-solid	
	phase	e extraction. The ann	ealed RNA duplex is further ana	lyzed by mass	
	spect	rometry to verify the	exact composition of the duple	ex. Each lot is compared to	
	the pi	revious lot by mass s	pectrometry to ensure maximu	m lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	e MARCH2 gene. Ead	h vial contains 5 nmol of lyophi	lized siRNA. The duplexes	
	can b	e transfected individ	ually or pooled together to achi	eve knockdown of the	
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	MAR	CH2 siRNA (Mouse)	- A 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

MARCH2 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
MARCH2 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
_		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 µl	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com