

Product Data Sheet

EPB41L5 siRNA (Mouse)

e Reactivity	Applications	
etic M	RNAi	
siRNA to inhibit EPB41L5 expression using RNA interference		
EPB41L5 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed		
to knock down gene expression.		
Lyophilized powder		
EPB41L5		
EPB4.1L5; KIAA1548; Band 4.1-like protein 5		
226352 (Mouse)		
Q8BGS1 (Mouse)		
> 97%		
Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure		
appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid		
phase extraction. The annealed RNA duplex is further analyzed by mass		
spectrometry to verify the exact composition of the duplex. Each lot is compared to		
the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.		
We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of		
mouse EPB41L5 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes		
can be transfected individually or po	ooled together to achieve	e knockdown of the
target gene, which is most commonly assessed by qPCR or western blot. Component 15 nmol 30 nmol		
EPB41L5 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
	etic M siRNA to inhibit EPB41L5 expression EPB41L5 siRNA (Mouse) is a target-site to knock down gene expression. Lyophilized powder EPB41L5 EPB41L5; KIAA1548; Band 4.1-like pi 226352 (Mouse) Q8BGS1 (Mouse) > 97% Oligonucleotide synthesis is monitor appropriate coupling efficiency. The phase extraction. The annealed RNA spectrometry to verify the exact corr the previous lot by mass spectrome We offers pre-designed sets of 3 diff mouse EPB41L5 gene. Each vial corr can be transfected individually or po target gene, which is most common Component EPB41L5 siRNA (Mouse) - A	tic M RNAi siRNA to inhibit EPB41L5 expression using RNA interference EPB41L5 siRNA (Mouse) is a target-specific 19-23 nt siRNA o to knock down gene expression. Lyophilized powder EPB41L5 EPB4.1L5; KIAA1548; Band 4.1-like protein 5 226352 (Mouse) Q8BGS1 (Mouse) > 97% Oligonucleotide synthesis is monitored base by base through appropriate coupling efficiency. The oligo is subsequently purphase extraction. The annealed RNA duplex is further analyze spectrometry to verify the exact composition of the duplex. the previous lot by mass spectrometry to ensure maximum I We offers pre-designed sets of 3 different target-specific siRI mouse EPB41L5 gene. Each vial contains 5 nmol of lyophilized can be transfected individually or pooled together to achieve target gene, which is most commonly assessed by qPCR or w Component 15 nmol EPB41L5 siRNA (Mouse) - A 5 nmol x 1

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

EP	B41L5 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Ne	gative Control	2.5 nmol x 1	2.5 nmol x 2
DE	PC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
of medium	of siRNA		2000
	100 nM	0.5 μl	0.25 μl
100 µl	50 nM	0.25 μl	0.25 μl
	10 nM	0.05 μl	0.25 μl
	100 nM	2.5 μl	1 µl
500 μl	50 nM	1.25 μl	1 µl
	10 nM	0.25 μl	1 µl
	100 nM	5 μl	2 µl
1 ml	50 nM	2.5 μl	2 µl
	10 nM	0.5 μl	2 µl
	100 nM	10 µl	5 µl
2 ml	50 nM	5 μl	5 μl
	10 nM	1 µl	5 μl
	of medium 100 μl 500 μl 1 ml	of medium of siRNA 100 nM 100 nM 100 nM 10 nM 50 nM 10 nM 500 μl 50 nM 100 nM 10 nM 500 μl 50 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 50 nM 1 nn 50 nM 10 nM 10 nM 10 nM 50 nM	of mediumof siRNA100 nM0.5 μl100 μl50 nM0.25 μl10 nM0.05 μl500 μl10 nM2.5 μl500 μl50 nM1.25 μl10 nM0.25 μl10 nM0.25 μl10 nM0.25 μl100 nM5 μl100 nM5 μl100 nM10 μl100 nM5.0 μl100 nM5.0 μl100 nM5.0 μl100 nM5.0 μl100 nM5.0 μl100 nM5.0 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com