

# **Product Data Sheet**

## CDC42BPA siRNA (Mouse)

Catalog #	Source	Reactivity	Applications		
CRN2538	Synthetic	Μ	RNAi		
Description	siRNA	to inhibit CDC42BPA	expression using RNA interfer	ence	
Specificity	CDC4	2BPA siRNA (Mouse)	is a target-specific 19-23 nt siR	NA oligo duplexes designed	
	to kno	ock down gene expre	ssion.		
Form	Lyoph	ilized powder			
Gene Symbol	CDC4	CDC42BPA			
Alternative N	ames KIAAC	KIAA0451; Serine/threonine-protein kinase MRCK alpha; CDC42-binding protein			
	kinase	e alpha			
Entrez Gene	22675	51 (Mouse)			
SwissProt	Q3UU	Q3UU96 (Mouse)			
Purity	> 97%	> 97%			
Quality Control Oligonucleotide synthesis is			s monitored base by base thro	ough trityl analysis to ensure	
	appro	priate coupling effici	ency. The oligo is subsequently	v purified by affinity-solid	
	phase	e extraction. The anne	ealed RNA duplex is further and	alyzed by mass	
	spect	rometry to verify the	exact composition of the dupl	ex. Each lot is compared to	
	the pi	revious lot by mass sp	pectrometry to ensure maximu	ım lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	e CDC42BPA gene. Ea	ch vial contains 5 nmol of lyop	hilized siRNA. The duplexes	
	can b	e transfected individu	ally or pooled together to ach	ieve knockdown of the	
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	CDC	42BPA siRNA (Mouse	) - A 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

CDC42BPA siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
CDC42BPA siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
_		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 μΙ
		10 nM	1 μΙ	5 μΙ

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com