

Product Data Sheet

TMEM194B siRNA (Mouse)

Catalog #	Source	Reactivity	Appl	ications		
CRN2559	Synthetic	Μ	RNAi			
Description	siRNA	to inhibit TMEM194B	expression using RN	A interference		
Specificity	TMEN	TMEM194B siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes				
	desigr	ned to knock down ger	e expression.			
Form	Lyoph	ilized powder				
Gene Symbol	TMEM	TMEM194B				
Alternative Na	ames Transr	Transmembrane protein 194B				
Entrez Gene	22709	94 (Mouse)				
SwissProt	Q8CB	Q8CB65 (Mouse)				
Purity	> 97%	> 97%				
Quality Contro	ality Control Oligonucleotide synthesis is monitored base by base through trityl analysis			base through trityl analysis to ensure		
	appro	priate coupling efficie	ncy. The oligo is subs	equently purified by affinity-solid		
	phase	extraction. The annea	led RNA duplex is fu	rther analyzed by mass		
	spectr	rometry to verify the e	xact composition of t	the duplex. Each lot is compared to		
	the pr	evious lot by mass spe	ectrometry to ensure	maximum lot-to-lot consistency.		
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	mouse	mouse TMEM194B gene. Each vial contains 5 nmol of lyophilized siRNA. The				
	duple	duplexes can be transfected individually or pooled together to achieve knockdown				
	of the	of the target gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nm	ol 30 nmol		
	TME	M194B siRNA (Mouse)	- A 5 nmo	l x 1 5 nmol x 2		
	TME	M194B siRNA (Mouse)	- B 5 nmo	l x 1 5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com