

Product Data Sheet

ZER1 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | |
|--|-------------|---|---------------------------------|-------------------------|--|
| CRN2607 | Synthetic | Μ | RNAi | | |
| Description siRNA | | o inhibit ZER1 expression using RNA interference | | | |
| Specificity | ZER1 s | ZER1 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| kno | | knock down gene expression. | | | |
| Form | Lyophi | Lyophilized powder | | | |
| Gene Symbol | ZER1 | ZER1 | | | |
| Alternative N | ames ZYG; Z | ZYG; ZYG11BL; Protein zer-1 homolog; Zyg-11 homolog B-like protein | | | |
| Entrez Gene | 22769 | 227693 (Mouse) | | | |
| SwissProt | Q80ZJ | Q80ZJ6 (Mouse) | | | |
| Purity | > 97% | > 97% | | | |
| Quality Contr | ol Oligon | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | |
| | approp | appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid | | | |
| | phase | phase extraction. The annealed RNA duplex is further analyzed by mass | | | |
| | spectro | spectrometry to verify the exact composition of the duplex. Each lot is compared to | | | |
| | the pre | evious lot by mass spec | trometry to ensure maximum | lot-to-lot consistency. | |
| Components | We off | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | mouse | ZER1 gene. Each vial co | ontains 5 nmol of lyophilized s | siRNA. The duplexes can | |
| | be trar | nsfected individually or | pooled together to achieve kr | nockdown of the target | |
| gene, which is most commonly assessed by qPCR or western blot. | | | n blot. | | |
| | Comp | oonent | 15 nmol | 30 nmol | |
| | ZER1 | siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

5 nmol x 1

5 nmol x 2

ZER1 siRNA (Mouse) - B

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| DEPC Water | 1 ml x 1 | 1 ml x 2 |
|------------------------|--------------|--------------|
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| ZER1 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 µl |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 μl | 2 µl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 µl |
| | | 10 nM | 0.5 μl | 2 µl |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 µl |
| | | 10 nM | 1 µl | 5 µl |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |