

Product Data Sheet

PPP1R37 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | |
|---|------------|--|-------------------------------------|---------------------------|--|
| CRN3034 | Synthetic | Μ | RNAi | | |
| Description | siRNA | to inhibit PPP1R37 e | xpression using RNA interference | е | |
| Specificity | PPP1F | PPP1R37 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed | | | |
| | to kno | ock down gene expre | ssion. | | |
| Form | Lyoph | ilized powder | | | |
| Gene Symbol | PPP1F | PPP1R37 | | | |
| Alternative N | ames LRRC6 | LRRC68; Protein phosphatase 1 regulatory subunit 37; Leucine-rich | | | |
| | repea | t-containing protein (| 58 | | |
| Entrez Gene | 23294 | 17 (Mouse) | | | |
| SwissProt Q | | Q8BKR5 (Mouse) | | | |
| Purity > 97% | | | | | |
| Quality Control Oligonucleotide synthesis is monitored base by base through trityl analys | | | gh trityl analysis to ensure | | |
| | appro | priate coupling efficion | ency. The oligo is subsequently p | urified by affinity-solid | |
| | phase | phase extraction. The annealed RNA duplex is further analyzed by mass | | | |
| | spect | rometry to verify the | exact composition of the duplex | . Each lot is compared to | |
| | the pi | revious lot by mass sp | ectrometry to ensure maximum | lot-to-lot consistency. | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | mous | e PPP1R37 gene. Eacl | n vial contains 5 nmol of lyophiliz | zed siRNA. The duplexes | |
| | can b | e transfected individu | ally or pooled together to achiev | ve knockdown of the | |
| | target | target gene, which is most commonly assessed by qPCR or western blot. | | | |
| | Com | ponent | 15 nmol | 30 nmol | |
| | PPP1 | LR37 siRNA (Mouse) - | A 5 nmol x 1 | 5 nmol x 2 | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| PPP1R37 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
|---------------------------|--------------|--------------|
| PPP1R37 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 µl |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 μl | 2 µl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 µl |
| | | 10 nM | 0.5 μl | 2 μl |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 µl | 5 μl |
| | | 10 nM | 1 μl | 5 μl |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |