

Product Data Sheet

C1QTNF5 siRNA (Mouse)

Catalog #	Source	Reactivity		Applications	
CRN3221	Synthetic	М		RNAi	
Description	siRNA	to inhibit C1QTNF5	expression usi	ng RNA interference	
Specificity	C1QT	NF5 siRNA (Mouse) i	s a target-spec	fic 19-23 nt siRNA oli	igo duplexes designed
	to kno	ock down gene expre	ession.		
Form	Lyoph	ilized powder			
Gene Symbol	C1QT	NF5			
Alternative N	ames CTRP	5; Complement C1q	tumor necrosis	factor-related protei	n 5
Entrez Gene	23532	12 (Mouse)			
SwissProt	Q8K4	79 (Mouse)			
Purity	> 97%	, D			
Quality Control Oligonucleotide synthesis is monitored base by base through trityl analy			trityl analysis to ensure		
	appro	priate coupling effic	iency. The oligo	is subsequently puri	ified by affinity-solid
	phase	e extraction. The ann	ealed RNA dup	lex is further analyze	d by mass
	spect	rometry to verify the	e exact compos	ition of the duplex. Ea	ach lot is compared to
	the p	revious lot by mass s	pectrometry to	ensure maximum lo	t-to-lot consistency.
Components	We of	ffers pre-designed se	ts of 3 differen	t target-specific siRN	A oligo duplexes of
	mous	e C1QTNF5 gene. Ea	ch vial contains	5 nmol of lyophilized	d siRNA. The duplexes
	can b	e transfected individ	ually or pooled	together to achieve	knockdown of the
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent		15 nmol	30 nmol
	C1Q	TNF5 siRNA (Mouse)	- A	5 nmol x 1	5 nmol x 2
	C1Q	TNF5 siRNA (Mouse)	- B	5 nmol x 1	5 nmol x 2

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

Negative Control 2.5 nmol x 1 2.5 nmol x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com