

# **Product Data Sheet**

## TRP53I11 siRNA (Mouse)

Catalog #	Source	Reactivity	Applica	ations	
CRN4199	Synthetic	М	RNAi		
Description	siRNA	to inhibit TRP53I11	expression using RNA int	erference	
Specificity	TRP53	3I11 siRNA (Mouse) i	a target-specific 19-23 r	nt siRNA oligo duplexes designed	
	to kno	ock down gene expre	ssion.		
Form	Lyoph	ilized powder			
Gene Symbol	TRP53	3111			
Alternative N	ames PIG11	PIG11; TP53I11; Tumor protein p53-inducible protein 11; Transformation related			
	protei	in 53 inducible prote	n 11; p53-induced gene	11 protein	
Entrez Gene	27741	L4 (Mouse)			
SwissProt	Q4QC	M4 (Mouse)			
Purity	> 97%				
Quality Contr	ol Oligor	nucleotide synthesis	s monitored base by bas	e through trityl analysis to ensure	
	appro	priate coupling effici	ency. The oligo is subseq	uently purified by affinity-solid	
	phase	extraction. The ann	ealed RNA duplex is furth	er analyzed by mass	
	spect	rometry to verify the	exact composition of the	e duplex. Each lot is compared to	
	the pr	revious lot by mass s	pectrometry to ensure m	aximum lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	e TRP53I11 gene. Ead	h vial contains 5 nmol of	lyophilized siRNA. The duplexes	
	can be	e transfected individ	ally or pooled together	to achieve knockdown of the	
	target	gene, which is most	commonly assessed by a	qPCR or western blot.	
	Com	ponent	15 nmol	30 nmol	
	TRP5	53I11 siRNA (Mouse)	- A 5 nmol x	1 5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

TRP53I11 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
TRP53I11 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 μl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 μl
		10 nM	1 μl	5 μl

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com