

Product Data Sheet

TAS2R136 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | |
|--|-----------|---|-------------------------------------|-----------------------------|--|
| CRN4736 | Synthetic | Μ | RNAi | | |
| Description | siRNA | A to inhibit TAS2R136 | expression using RNA interfere | nce | |
| Specificity | TAS2 | R136 siRNA (Mouse) | s a target-specific 19-23 nt siRN | IA oligo duplexes designed | |
| | to kn | ock down gene expre | ssion. | | |
| Form | Lyoph | nilized powder | | | |
| Gene Symbol | TAS2 | TAS2R136 | | | |
| Alternative N | ames T2R5 | T2R52; TAS2R36; Taste receptor type 2 member 136; T2R136; Taste receptor type 2 | | | |
| | mem | ber 36; T2R36; mT2r! | 52 | | |
| Entrez Gene | 3531 | 65 (Mouse) | | | |
| SwissProt | Q7TC | Q7TQA8 (Mouse) | | | |
| Purity | > 97% | > 97% | | | |
| Quality Control O | | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | |
| | appro | opriate coupling effici | ency. The oligo is subsequently | purified by affinity-solid | |
| | phase | e extraction. The ann | ealed RNA duplex is further ana | lyzed by mass | |
| | spect | rometry to verify the | exact composition of the duple | ex. Each lot is compared to | |
| | the p | revious lot by mass s | pectrometry to ensure maximu | m lot-to-lot consistency. | |
| Components We offers pre-designed sets of 3 diffe | | | ts of 3 different target-specific s | siRNA oligo duplexes of | |
| | mous | se TAS2R136 gene. Ea | ch vial contains 5 nmol of lyoph | ilized siRNA. The duplexes | |
| | can b | e transfected individ | ually or pooled together to achi | eve knockdown of the | |
| | targe | target gene, which is most commonly assessed by qPCR or western blot. | | | |
| | Com | ponent | 15 nmol | 30 nmol | |
| | TAS2 | 2R136 siRNA (Mouse) | - A 5 nmol x 1 | 5 nmol x 2 | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| TAS2R136 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
|----------------------------|--------------|--------------|
| TAS2R136 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μl |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 μl | 2 µl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 µl |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μΙ |
| | | 10 nM | 1 μl | 5 μΙ |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |