

For research purposes only, not for human use

Product Data Sheet

Recombinant Human HDGF Protein

| Catalog # | Source | Reactivity | Applications | | | |
|--------------------|---------|--|---|--|--|--|
| CRP1662 | E. coli | Human | E, WB, SDS-PAGE, MS | | | |
| Description | | Recombinant Human HDGF | Protein is produced by our E. coli expression system and | | | |
| | | the target gene encoding N | let1-Tyr100 is expressed with a 6His tag at the | | | |
| | | C-terminus. | | | | |
| Form | | Lyophilized from a 0.2 μ M f | iltered solution of 20mM Tris, 1mM DTT, 1mM EDTA, pH | | | |
| | | 7.5. | | | | |
| Gene Symbol | | HDGF | | | | |
| Alternative Names | | HMG1L2; Hepatoma-derived growth factor; HDGF; High mobility group protein | | | | |
| | | 1-like 2; HMG-1L2 | | | | |
| Entrez Gene | | 3068 (Human) | | | | |
| SwissProt | | P51858 (Human) | | | | |
| Purity | | Greater than 95% as determined by reducing SDS-PAGE. | | | | |
| Chemical Structure | | MSRSNRQKEY KCGDLVFAKM KGYPHWPARI DEMPEAAVKS TANKYQVFFF GTHETAFLGP | | | | |
| | | KDLFPYEESK EKFGKPNKRK | GFSEGLWEIE NNPTVKASGY LEHHHHHH | | | |
| Quality Contr | ol | Endotoxin: Less than 0.1 ng | /μg (1 IEU/μg) as determined by LAL test. | | | |
| Directions for | r Use | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not | | | | |
| | | recommended to reconstit | ute to a concentration less than 100 μ g/ml. Dissolve the | | | |
| | | lyophilized protein in 1X PB | S. Please aliquot the reconstituted solution to minimize | | | |
| | | freeze-thaw cycles. | | | | |
| Storage/Stab | ility | Lyophilized protein should | be stored at -20°C, though stable at room temperature | | | |
| | | for 3 weeks. Reconstituted | protein solution can be stored at 2-8°C for 2-7 days. | | | |
| | | Aliquots of reconstituted sa | imples are stable at -20°C for 3 months. | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

Product Data Sheet

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com