

## Recombinant Human NHP2L1 Protein

| Catalog #                 | Source   | Reactivity | Applications        |
|---------------------------|--|------------|---------------------|
| CRP2012                   | E. coli  | Human      | E, WB, SDS-PAGE, MS |
| <b>Description</b>        | Recombinant Human NHP2L1 Protein is produced by our E.coli expression system and the target gene encoding Met1-Val128 is expressed with a 6His tag at the N-terminus.  |            |                     |
| <b>Form</b>               | Lyophilized from a 0.2 μM filtered solution of 20mM Tris-HCl, 600mM NaCl, pH 8.0.  |            |                     |
| <b>Gene Symbol</b>        | NHP2L1   |            |                     |
| <b>Alternative Names</b>  | NHP2-like protein 1; High mobility group-like nuclear protein 2 homolog 1; OTK27; SNU13 homolog; hSNU13; U4/U6.U5 tri-snRNP 15.5 kDa protein   |            |                     |
| <b>Entrez Gene</b>        | 4809 (Human)   |            |                     |
| <b>SwissProt</b>          | P55769 (Human)   |            |                     |
| <b>Purity</b>             | Greater than 95% as determined by reducing SDS-PAGE.   |            |                     |
| <b>Chemical Structure</b> | MGSSHHHHHH SSGLVPRGSH MTEADVNPKA YPLADAHLTK KLLDLVQQSC<br>NYKQLRKGAN EATKTLNRGI SEFIVMAADA EPLEIILHLP LLCEDKNVPY VFVRSKQALG<br>RACGVS RPVI ACSVTIKEGS QLKQQIQSIQ QSIERLLV  |            |                     |
| <b>Quality Control</b>    | Endotoxin: Less than 0.1 ng/μg (1 IEU/μg) as determined by LAL test.   |            |                     |
| <b>Directions for Use</b> | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/ml. Dissolve the lyophilized protein in 1X PBS. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |            |                     |
| <b>Storage/Stability</b>  | Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  |            |                     |

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, CHIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

### COHESION BIOSCIENCES LIMITED

**WEB**  
[www.cohesionbio.com](http://www.cohesionbio.com)

**ORDER**  
[order@cohesionbio.com](mailto:order@cohesionbio.com)

**SUPPORT**  
[techsupport@cohesionbio.com](mailto:techsupport@cohesionbio.com)

**CUSTOM**  
[custom@cohesionbio.com](mailto:custom@cohesionbio.com)

# Product Data Sheet

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, CHIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

## COHESION BIOSCIENCES LIMITED

**WEB**

[www.cohesionbio.com](http://www.cohesionbio.com)

**ORDER**

[order@cohesionbio.com](mailto:order@cohesionbio.com)

**SUPPORT**

[techsupport@cohesionbio.com](mailto:techsupport@cohesionbio.com)

**CUSTOM**

[custom@cohesionbio.com](mailto:custom@cohesionbio.com)