

## Recombinant Human IL-1R1 Protein

| Catalog #                 | Source   | Reactivity | Applications        |
|---------------------------|--|------------|---------------------|
| CRP2739                   | Human cells  | Human      | E, WB, SDS-PAGE, MS |
| <b>Description</b>        | Recombinant Human IL-1R1 Protein is produced by mammalian expression system and the target gene encoding Leu18-Thr332 is expressed fused with a 6His tag at the C-terminus.  |            |                     |
| <b>Form</b>               | Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.   |            |                     |
| <b>Gene Symbol</b>        | IL1R1  |            |                     |
| <b>Alternative Names</b>  | IL1R; IL1RA; IL1RT1; Interleukin-1 receptor type 1; IL-1R-1; IL-1RT-1; IL-1RT1; CD121 antigen-like family member A; Interleukin-1 receptor alpha; IL-1R-alpha; Interleukin-1 receptor type I; p80; CD121a  |            |                     |
| <b>Entrez Gene</b>        | 3554 (Human)   |            |                     |
| <b>SwissProt</b>          | P14778 (Human)   |            |                     |
| <b>Purity</b>             | Greater than 95% as determined by reducing SDS-PAGE.   |            |                     |
| <b>Chemical Structure</b> | LEADKCKERE EKIILVSSAN EIDVRPCPLN PNEHKGITITW YKDDSKTPVS TEQASRIHQH<br>KEKLWFVPAK VEDSGHYICV VRNSSYCLRI KISAKFVENE PNLCYNAQAI FKQKLPVAGD<br>GGLVCPYMEF FKNENNELPK LQWYKDCCKPL LLDNIHFSGV KDRLIVMNVA EKHRGNYTCH<br>ASYTYLGKQY PITRVIEFIT LEENKPTRPV IVSPANETME VDLGSQIQLI CNVTGQLSDI<br>AYWKWNGSVI DEDDPVLGED YYSVENPANK RRSTLITVLN ISEIESRFYK HPFTCFKANT<br>HGIDAAYIQL IYPVTHHHHH H |            |                     |
| <b>Quality Control</b>    | Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.   |            |                     |
| <b>Directions for Use</b> | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize  |            |                     |

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

### COHESION BIOSCIENCES LIMITED

**WEB**

[www.cohesionbio.com](http://www.cohesionbio.com)

**ORDER**

[order@cohesionbio.com](mailto:order@cohesionbio.com)

**SUPPORT**

[techsupport@cohesionbio.com](mailto:techsupport@cohesionbio.com)

**CUSTOM**

[custom@cohesionbio.com](mailto:custom@cohesionbio.com)

# Product Data Sheet

freeze-thaw cycles.

## Storage/Stability

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 2-8°C for 2-7 days.

Aliquots of reconstituted samples are stable at -20°C for 3 months.

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, CHIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

## COHESION BIOSCIENCES LIMITED

### WEB

[www.cohesionbio.com](http://www.cohesionbio.com)

### ORDER

[order@cohesionbio.com](mailto:order@cohesionbio.com)

### SUPPORT

[techsupport@cohesionbio.com](mailto:techsupport@cohesionbio.com)

### CUSTOM

[custom@cohesionbio.com](mailto:custom@cohesionbio.com)