

Bovine IgG

Catalog #	Source	Reactivity	Applications
CRP9000	Bovine		WB, IH, IP

Description The purified total Bovine IgG can be used as a negative control for antibodies in Western blot, immunoprecipitation, and immunohistochemistry, etc.

Form Purified Bovine IgG in PBS, pH 7.3, 0.01% sodium azide.

Gene Symbol

Alternative Names

Entrez Gene

SwissProt

Purity Greater than 95% as determined by reducing SDS-PAGE.

Chemical Structure

Quality Control

Directions for Use Please minimize freeze-thaw cycles.

Storage/Stability Store it at -20°C to -80°C for one year. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB

www.cohesionbio.com

ORDER

order@cohesionbio.com

SUPPORT

techsupport@cohesionbio.com

CUSTOM

custom@cohesionbio.com