

Product Data Sheet

CTH siRNA (Rat)

Catalog # Sourc	e Reactivity	Applications		
CRR0384 Synth	etic R	RNAi		
Description	Description siRNA to inhibit CTH expression using RNA interference			
Specificity	CTH siRNA (Rat) is a target-specific 19-23 nt siRNA oligo duplexes designed to knock			
	down gene expression.			
Form	Lyophilized powder			
Gene Symbol CTH				
Alternative Names Cystathionine gamma-lyase; Cysteine-protein sulfhydrase; Gamma-cystath		mma-cystathionase;		
	Probasin-related antigen; P	RB-RA		
Entrez Gene	ntrez Gene 24962 (Rat)			
SwissProt	P18757 (Rat)			
Purity	> 97%			
Quality Control	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid			
	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.			
Components	mponents We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of rat			
	CTH gene. Each vial contain	is 5 nmol of lyophilized siRNA. The	duplexes can be	
	transfected individually or	pooled together to achieve knockdo	own of the target	
	gene, which is most commonly assessed by qPCR or western blot.			
	Component	15 nmol	30 nmol	
	CTH siRNA (Rat) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

CTH siRNA (Rat) - B	5 nmol x 1	5 nmol x 2
CTH siRNA (Rat) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 μΙ
		10 nM	1 μΙ	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com