

# **Product Data Sheet**

### CSRP2 siRNA (Rat)

Catalog #	Source	Reactivity	Applications			
CRR1074	Synthetic	R	RNAi			
Description	siRNA	to inhibit CSRP2 expr	ession using RNA interference			
Specificity	CSRP2	CSRP2 siRNA (Rat) is a target-specific 19-23 nt siRNA oligo duplexes designed to				
	knock	down gene expressio	n.			
Form	Lyoph	ilized powder				
Gene Symbol	CSRP2	CSRP2				
Alternative N	ames SMLIN	SMLIM; Cysteine and glycine-rich protein 2; Cysteine-rich protein 2; CRP2; Smooth				
	muscl	e cell LIM protein; Sm	LIM			
Entrez Gene	29317	(Rat)				
SwissProt	Q6290	Q62908 (Rat)				
Purity > 9		> 97%				
Quality Control Oligonucleotide synthesis is monitored base by base th			monitored base by base through	ugh trityl analysis to ensure		
appropriate coupling efficiency. The oligo			ncy. The oligo is subsequently pu	is subsequently purified by affinity-solid		
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass				
	spectr	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the pr	the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.				
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of rat				
	CSRP2	CSRP2 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can be				
	transf	transfected individually or pooled together to achieve knockdown of the target				
	gene,	gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
	CSRP	2 siRNA (Rat) - A	5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

CSRP2 siRNA (Rat) - B	5 nmol x 1	5 nmol x 2
CSRP2 siRNA (Rat) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 μl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
_		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 µl	5 μΙ

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com