

Product Data Sheet

ARHGAP17 siRNA (Rat)

Catalog #	Source	Reactivity	Applications			
CRR1773	Synthetic	R	RNAi			
Description	siRNA	to inhibit ARHGAP17	expression using RNA interference	ce		
Specificity	ARHG	ARHGAP17 siRNA (Rat) is a target-specific 19-23 nt siRNA oligo duplexes designed to				
	knock	down gene expressio	on.			
Form	Lyoph	ilized powder				
Gene Symbol	ARHG	ARHGAP17				
Alternative N	ames Rho G	Rho GTPase-activating protein 17; Neuron-associated developmentally-regulated				
	protei	in; Nadrin; Rho-type (GTPase-activating protein 17			
Entrez Gene	63994	63994 (Rat)				
SwissProt	Q99N	Q99N37 (Rat)				
Purity	> 97%	> 97%				
Quality Control Oligonucleotide synthesis is monitored base by base through trityl analy			n trityl analysis to ensure			
	appro	priate coupling efficie	ency. The oligo is subsequently pu	rified by affinity-solid		
	phase	e extraction. The anne	aled RNA duplex is further analyz	ed by mass		
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the pr	revious lot by mass sp	ectrometry to ensure maximum l	ot-to-lot consistency.		
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of rat				
	ARHG	ARHGAP17 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can be				
	transf	transfected individually or pooled together to achieve knockdown of the target				
	gene,	gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
	ARH	GAP17 siRNA (Rat) - A	5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

ARHGAP17 siRNA (Rat) - B	5 nmol x 1	5 nmol x 2
ARHGAP17 siRNA (Rat) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μl
_		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 μΙ
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com