

Product Data Sheet

CDK5RAP3 siRNA (Rat)

Catalog #	Source	Reactivity	Applications		
CRR2215	Synthetic	R	RNAi		
Description si		siRNA to inhibit CDK5RAP3 expression using RNA interference			
Specificity	CDK5	CDK5RAP3 siRNA (Rat) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knock	k down gene expressi	on.		
Form	Lyoph	nilized powder			
Gene Symbol	CDK5	CDK5RAP3			
Alternative N	ames CDK5	CDK5 regulatory subunit-associated protein 3; CDK5 activator-binding protein C53			
Entrez Gene	80278	80278 (Rat)			
SwissProt	Q9JLH	Q9JLH7 (Rat)			
Purity >		> 97%			
Quality Control Oligonucleotide synthesis is monitored base by base through tri-			h trityl analysis to ensure		
	appro	opriate coupling effici	ency. The oligo is subsequently pu	urified by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the p	revious lot by mass s	pectrometry to ensure maximum	lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of rat			
	CDK5	CDK5RAP3 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can be			
	transf	transfected individually or pooled together to achieve knockdown of the target			
	gene,	gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	CDK	5RAP3 siRNA (Rat) - A	5 nmol x 1	5 nmol x 2	

CDK5RAP3 siRNA (Rat) - B5 nmol x 15 nmol x 2Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-
Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility
Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference
Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-

Rabbit, S- Sheep, Z- Zebrafish COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	СИЗТОМ
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

CDK5RAP3 siRNA (Rat) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 μl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com