

Product Data Sheet

ACTR3 siRNA (Rat)

	•				
Catalog #	Source	Reactivity	Applications		
CRR2326	Synthetic	R	RNAi		
Description	Description siRNA to inhibit ACTR3 expression using RNA interference				
Specificity	ACTR	ACTR3 siRNA (Rat) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knock	down gene expression	on.		
Form	Lyoph	Lyophilized powder			
Gene Symbol	ACTR	ACTR3			
Alternative N	ames ARP3;	ARP3; Actin-related protein 3; Actin-like protein 3			
Entrez Gene	81732	81732 (Rat)			
SwissProt	Q4V7	Q4V7C7 (Rat)			
Purity	> 97%	> 97%			
Quality Control Oligonucleotide synthesis is monitored base by base through trityl anal			ugh trityl analysis to ensure		
appropriate coupling efficiency. The oligo i		ency. The oligo is subsequently	is subsequently purified by affinity-solid		
	phase	e extraction. The anne	ealed RNA duplex is further and	alyzed by mass	
	spect	rometry to verify the	exact composition of the duple	ex. Each lot is compared to	
	the p	revious lot by mass sp	pectrometry to ensure maximu	m lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of rat			
	ACTR	3 gene. Each vial cont	ains 5 nmol of lyophilized siRN	A. The duplexes can be	
	transf	fected individually or	pooled together to achieve kno	ockdown of the target	
	gene, which is most commonly assessed by qPCR or western blot.			ern blot.	
	Com	ponent	15 nmol	30 nmol	
	ACTI	R3 siRNA (Rat) - A	5 nmol x 1	5 nmol x 2	
	ACTI	R3 siRNA (Rat) - B	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

ACTR3 siRNA (Rat) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com