

Product Data Sheet

CIB1 siRNA (Rat)

| Catalog # | Source | Reactivity | Applications | | |
|---------------------|--------------------------------------------------------------------------------|----------------------------------------------------------------------------------------|-----------------------------------------------------|------------------------|--|
| CRR2391 | Synthetic | R | RNAi | | |
| Description | siRNA | to inhibit CIB1 expre | ession using RNA interference | | |
| Specificity | CIB1 s | siRNA (Rat) is a target | -specific 19-23 nt siRNA oligo duple | exes designed to knock | |
| | down | gene expression. | | | |
| Form | Lyoph | nilized powder | | | |
| Gene Symbol | CIB1 | CIB1 | | | |
| Alternative N | ames CIB; K | CIB; KIP; PRKDCIP; Calcium and integrin-binding protein 1; CIB; Calmyrin; | | | |
| | DNA- | PKcs-interacting prote | ein; Kinase-interacting protein; KIP | | |
| Entrez Gene | 81823 | 3 (Rat) | | | |
| SwissProt | Q9R0 | Q9R010 (Rat) | | | |
| Purity | > 97% | > 97% | | | |
| Quality Control Oli | | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | |
| | appropriate coupling efficiency. The oligo is subsequently purified by affinit | | | fied by affinity-solid | |
| phase extraction. | | e extraction. The anne | The annealed RNA duplex is further analyzed by mass | | |
| | spect | spectrometry to verify the exact composition of the duplex. Each lot is compared to | | | |
| | the p | revious lot by mass sp | pectrometry to ensure maximum lo | t-to-lot consistency. | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of rat | | | |
| | CIB1 g | CIB1 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can be | | | |
| | transf | transfected individually or pooled together to achieve knockdown of the target | | | |
| | gene, | gene, which is most commonly assessed by qPCR or western blot. | | | |
| | Com | ponent | 15 nmol | 30 nmol | |
| | CIB1 | siRNA (Rat) - A | 5 nmol x 1 | 5 nmol x 2 | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| CIB1 siRNA (Rat) - B | 5 nmol x 1 | 5 nmol x 2 |
|----------------------|--------------|--------------|
| CIB1 siRNA (Rat) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| _ | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 μl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 µl |
| | | 10 nM | 1 μΙ | 5 μΙ |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |