

Product Data Sheet

IDI1 siRNA (Rat)

Catalog # Sourc	e Reactivity	Applications		
CRR2691 Synth	etic R	RNAi		
Description	siRNA to inhibit IDI1 expres	ssion using RNA interference		
Specificity	IDI1 siRNA (Rat) is a target-specific 19-23 nt siRNA oligo duplexes designed to knock			
	down gene expression.			
Form	Lyophilized powder			
Gene Symbol	IDI1			
Alternative Names	Isopentenyl-diphosphate Delta-isomerase 1; Isopentenyl pyrophosphate isomerase			
	1; IPP isomerase 1; IPPI1			
Entrez Gene	89784 (Rat)			
SwissProt	O35760 (Rat)			
Purity	> 97%			
Quality Control	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensu			
	appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass			
spectrometry to verify the exact composition of the duplex. Each lot is compar			ch lot is compared to	
	the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.			
Components	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of rat			
	IDI1 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can be			
	transfected individually or pooled together to achieve knockdown of the target			
	gene, which is most commonly assessed by qPCR or western blot.			
	Component	15 nmol	30 nmol	
	IDI1 siRNA (Rat) - A	5 nmol x 1	5 nmol x 2	
Gene Symbol Alternative Names Entrez Gene SwissProt Purity Quality Control	IDI1 Isopentenyl-diphosphate D 1; IPP isomerase 1; IPPI1 89784 (Rat) O35760 (Rat) > 97% Oligonucleotide synthesis is appropriate coupling efficie phase extraction. The anne spectrometry to verify the the previous lot by mass sp We offers pre-designed set IDI1 gene. Each vial contain transfected individually or p gene, which is most common Component	s monitored base by base through t ency. The oligo is subsequently purif ealed RNA duplex is further analyzed exact composition of the duplex. Ea bectrometry to ensure maximum lot s of 3 different target-specific siRNA hs 5 nmol of lyophilized siRNA. The o pooled together to achieve knockdo only assessed by qPCR or western b 15 nmol	rityl analysis to ensur fied by affinity-solid d by mass ach lot is compared to to-to-lot consistency. A oligo duplexes of rat duplexes can be own of the target lot. 30 nmol	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

IDI1 siRNA (Rat) - B	5 nmol x 1	5 nmol x 2
IDI1 siRNA (Rat) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well 5	500 µl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
6-well	2 ml	100 nM	10 µl	5 μΙ
		50 nM	5 µl	5 μΙ
		10 nM	1 μΙ	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com