

Product Data Sheet

CX3CL1 siRNA (Rat)

Catalog #	Source	Reactivity	Applications			
CRR2699	Synthetic	R	RNAi			
Description	siRNA	to inhibit CX3CL1 ex	pression using RNA interference			
Specificity	CX3CL	1 siRNA (Rat) is a tar	get-specific 19-23 nt siRNA oligo du	plexes designed to		
	knock	down gene expression	on.			
Form	Lyoph	ilized powder				
Gene Symbol	CX3CL	CX3CL1				
Alternative N	ames ACC1;	ACC1; FKN; SCYD1; Fractalkine; C-X3-C motif chemokine 1; CX3C				
	memb	orane-anchored chem	nokine; Neurotactin; Small-inducible	e cytokine D1		
Entrez Gene	89808	3 (Rat)				
SwissProt	05514	O55145 (Rat)				
Purity	> 97%	> 97%				
Quality Contr	ol Oligor	nucleotide synthesis i	eotide synthesis is monitored base by base through trityl analysis to ensure			
	appropriate coupling efficiency. The oligo is subsequently purified by affinity-			fied by affinity-solid		
phase extraction. The annealed RNA duplex is further analyzed b			d by mass			
	specti	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the pr	revious lot by mass sp	pectrometry to ensure maximum lo	t-to-lot consistency.		
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of rat				
	CX3CL	CX3CL1 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can be				
	transf	transfected individually or pooled together to achieve knockdown of the target				
	gene,	gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
	CX30	CL1 siRNA (Rat) - A	5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

CX3CL1 siRNA (Rat) - B	5 nmol x 1	5 nmol x 2
CX3CL1 siRNA (Rat) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 μl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 μl
		10 nM	1 μl	5 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com