

Product Data Sheet

AP2M1 siRNA (Rat)

| Catalog # | Source | Reactivity | Applications | | | |
|-----------------------|-----------|---|--|--|--|--|
| CRR2934 | Synthetic | R | RNAi | | | |
| Description | siRNA | to inhibit AP2M1 exp | pression using RNA interference | | | |
| Specificity | AP2N | 11 siRNA (Rat) is a targ | get-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| | knock | down gene expressio | n. | | | |
| Form | Lyoph | nilized powder | | | | |
| Gene Symbol | AP2N | 11 | | | | |
| Alternative Na | ames AP-2 | AP-2 complex subunit mu; AP-2 mu chain; Adapter-related protein complex 2 | | | | |
| | subur | nit mu; Adaptor protei | in complex AP-2 subunit mu; Clathrin assembly protein | | | |
| | comp | lex 2 mu medium cha | in; Clathrin coat assembly protein AP50; Clathrin | | | |
| | coat-a | associated protein AP | 50; | | | |
| Entrez Gene | 11656 | 63 (Rat) | | | | |
| SwissProt | P8409 | 92 (Rat) | | | | |
| Purity | > 97% | , 0 | | | | |
| Quality Contro | ol Oligoi | nucleotide synthesis is | s monitored base by base through trityl analysis to ensure | | | |
| | appro | priate coupling efficie | ency. The oligo is subsequently purified by affinity-solid | | | |
| | phase | extraction. The anne | aled RNA duplex is further analyzed by mass | | | |
| | spect | rometry to verify the | exact composition of the duplex. Each lot is compared to | | | |
| | the p | revious lot by mass sp | ectrometry to ensure maximum lot-to-lot consistency. | | | |
| Components | We of | ffers pre-designed set | s of 3 different target-specific siRNA oligo duplexes of rat | | | |
| | AP2N | 11 gene. Each vial con | tains 5 nmol of lyophilized siRNA. The duplexes can be | | | |
| | transf | fected individually or | pooled together to achieve knockdown of the target | | | |
| | gene, | which is most commo | only assessed by qPCR or western blot. | | | |
| A secold sectors from | | (anterna black III) terraria | histochemister, IC, January officerson and IC, Class externation, IC | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| Component | 15 nmol | 30 nmol |
|-----------------------|--------------|--------------|
| AP2M1 siRNA (Rat) - A | 5 nmol x 1 | 5 nmol x 2 |
| AP2M1 siRNA (Rat) - B | 5 nmol x 1 | 5 nmol x 2 |
| AP2M1 siRNA (Rat) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 μl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 µl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| 6-well | 2 ml | 100 nM | 10 µl | 5 µl |
| | | 50 nM | 5 µl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com