

Product Data Sheet

DCPS siRNA (Rat)

Source	Reactivity	Applications		
Synthetic	R	RNAi		
siRNA	to inhibit DCPS expre	ession using RNA interference		
DCPS	DCPS siRNA (Rat) is a target-specific 19-23 nt siRNA oligo duplexes designed to knock			
down	gene expression.			
Lyoph	nilized powder			
DCPS	DCPS			
mes DCS1	DCS1; HINT5; m7GpppX diphosphatase; DCS-1; Decapping scavenger enzyme;			
Hint-r	elated 7meGMP-dire	cted hydrolase; Histidine triad nucleotide-binding protein		
5; His	tidine triad protein m	ember 5; HINT-5; Scavenger mRNA-decapping enzyme		
DcpS				
26660	266605 (Rat)			
Q8K4	Q8K4F7 (Rat)			
> 97%	> 97%			
Oligo	nucleotide synthesis i	s monitored base by base through trityl analysis to ensure		
appro	opriate coupling efficie	ency. The oligo is subsequently purified by affinity-solid		
phase	e extraction. The anne	aled RNA duplex is further analyzed by mass		
spect	rometry to verify the	exact composition of the duplex. Each lot is compared to		
the p	revious lot by mass sp	ectrometry to ensure maximum lot-to-lot consistency.		
We o	ffers pre-designed set	s of 3 different target-specific siRNA oligo duplexes of rat		
DCPS	gene. Each vial conta	ins 5 nmol of lyophilized siRNA. The duplexes can be		
trans	fected individually or l	pooled together to achieve knockdown of the target		
gene,	which is most commo	only assessed by qPCR or western blot.		
	Synthetic siRNA DCPS down Lyoph DCPS DCS1 Hint-r 5; His DcpS 26660 Q8K4 > 97% Oligo appro phase spect the p We o DCPS	SyntheticRsiRNA to inhibit DCPS expressionDCPS siRNA (Rat) is a targetdown gene expression.Lyophilized powderDCPSmesDCS1; HINT5; m7GpppX digHint-related 7meGMP-dired5; Histidine triad protein mDcpS266605 (Rat)Q8K4F7 (Rat)> 97%Oligonucleotide synthesis isappropriate coupling efficiesphase extraction. The anneespectrometry to verify thethe previous lot by mass spWe offers pre-designed setDCPS gene. Each vial contaitransfected individually or		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

Component	15 nmol	30 nmol
DCPS siRNA (Rat) - A	5 nmol x 1	5 nmol x 2
DCPS siRNA (Rat) - B	5 nmol x 1	5 nmol x 2
DCPS siRNA (Rat) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μl
		100 nM	5 µl	2 μΙ
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
6-well	2	100 nM	10 µl	5 μl
	2 ml	50 nM	5 μl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com