

Product Data Sheet

CYP4F4 siRNA (Rat)

Catalog #	Source	Reactivity	Applications		
CRR3812	Synthetic	R	RNAi		
Description	siRNA	to inhibit CYP4F4 exp	ression using RNA interferenc	e	
Specificity	CYP4F	CYP4F4 siRNA (Rat) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knock	down gene expressio	n.		
Form	Lyoph	ilized powder			
Gene Symbol	CYP4F	CYP4F4			
Alternative N	ames Cytocl	Cytochrome P450 4F4; CYPIVF4			
Entrez Gene	28690	286904 (Rat)			
SwissProt	P5186	P51869 (Rat)			
Purity	> 97%	ı.			
Quality Control Oligonucleotide synthesis is monitored base			monitored base by base thro	ugh trityl analysis to ensure	
	appro	priate coupling efficie	ncy. The oligo is subsequently	purified by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spectr	ometry to verify the e	exact composition of the duple	ex. Each lot is compared to	
	the pr	evious lot by mass spe	ectrometry to ensure maximu	m lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of rat			
	CYP4F	CYP4F4 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can be			
	transf	transfected individually or pooled together to achieve knockdown of the target			
	gene,	gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	CYP4	F4 siRNA (Rat) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

5 nmol x 1

5 nmol x 2

CYP4F4 siRNA (Rat) - B

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

_	DEPC Water	1 ml x 1	1 ml x 2
	DEDC Mater	1 mal x 1	1 mal v 2
	Negative Control	2.5 nmol x 1	2.5 nmol x 2
	CYP4F4 siRNA (Rat) - C	5 nmol x 1	5 nmol x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com