

Product Data Sheet

PHAX siRNA (Rat)

| Catalog # | Source | Reactivity | Applications | | |
|-----------------------------------------------------------------------------------------|-----------|---------------------------------------------------------------------------------------|----------------------------------------------|---------------------------|--|
| CRR3820 | Synthetic | R | RNAi | | |
| Description | siRNA | to inhibit PHAX expre | ession using RNA interference | | |
| Specificity | РНАХ | PHAX siRNA (Rat) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| | knock | down gene expressio | n. | | |
| Form | Lyoph | ilized powder | | | |
| Gene Symbol | PHAX | РНАХ | | | |
| Alternative Na | ames RNUX | RNUXA; Phosphorylated adapter RNA export protein; 26 kDa resiniferatoxin-binding | | | |
| | protei | in; RBP-26; RNA U sma | all nuclear RNA export adapter p | protein; RTX-42; | |
| | Resini | feratoxin-binding prot | ein 2; RBP-2 | | |
| Entrez Gene | 28691 | .7 (Rat) | | | |
| SwissProt | Q6306 | Q63068 (Rat) | | | |
| Purity | > 97% | > 97% | | | |
| Quality Contro | ol Oligor | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | |
| | appro | priate coupling efficie | ncy. The oligo is subsequently p | urified by affinity-solid | |
| | phase | extraction. The annea | aled RNA duplex is further analy | zed by mass | |
| | spectr | rometry to verify the e | exact composition of the duplex. | . Each lot is compared to | |
| | the pr | evious lot by mass spe | ectrometry to ensure maximum | lot-to-lot consistency. | |
| Components We offers pre-designed sets of 3 different target-specific siRNA oligo duple | | | NA oligo duplexes of rat | | |
| | PHAX | gene. Each vial contai | ns 5 nmol of lyophilized siRNA. ⁻ | The duplexes can be | |
| | transf | ected individually or p | oooled together to achieve knoc | kdown of the target | |
| | gene, | which is most commo | only assessed by qPCR or wester | n blot. | |
| | Com | ponent | 15 nmol | 30 nmol | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| PHAX siRNA (Rat) - A | 5 nmol x 1 | 5 nmol x 2 |
|----------------------|--------------|--------------|
| PHAX siRNA (Rat) - B | 5 nmol x 1 | 5 nmol x 2 |
| PHAX siRNA (Rat) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 µl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 μl |
| 6-well | 2 ml | 50 nM | 5 µl | 5 µl |
| | | 10 nM | 1 μl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

Product Data Sheet

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com