

Product Data Sheet

METRN siRNA (Rat)

Reactivity	Applications				
tic R	RNAi				
iption siRNA to inhibit METRN expression using RNA interference					
METRN siRNA (Rat) is a target-spe	siRNA (Rat) is a target-specific 19-23 nt siRNA oligo duplexes designed to				
knock down gene expression.					
Lyophilized powder					
Gene Symbol METRN					
HYRAC; Meteorin; Hypoxia/reoxygenation regulatory factor					
287151 (Rat)					
Q5Q0T9 (Rat)					
> 97%					
Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure					
appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to					
			the previous lot by mass spectrom	netry to ensure maximum	lot-to-lot consistency.
			Components We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of rat METRN gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can be transfected individually or pooled together to achieve knockdown of the target gene, which is most commonly assessed by qPCR or western blot.		
Component	15 nmol	30 nmol			
METRN siRNA (Rat) - A	5 nmol x 1	5 nmol x 2			
METRN siRNA (Rat) - B	5 nmol x 1	5 nmol x 2			
	tic R siRNA to inhibit METRN expression METRN siRNA (Rat) is a target-spee knock down gene expression. Lyophilized powder METRN HYRAC; Meteorin; Hypoxia/reoxyg 287151 (Rat) Q5Q0T9 (Rat) > 97% Oligonucleotide synthesis is monit appropriate coupling efficiency. Th phase extraction. The annealed RI spectrometry to verify the exact of the previous lot by mass spectrom We offers pre-designed sets of 3 of METRN gene. Each vial contains 5 transfected individually or pooled gene, which is most commonly as: Component METRN siRNA (Rat) - A	tic R RNAi siRNA to inhibit METRN expression using RNA interference METRN siRNA (Rat) is a target-specific 19-23 nt siRNA oligo knock down gene expression. Lyophilized powder METRN HYRAC; Meteorin; Hypoxia/reoxygenation regulatory factor 287151 (Rat) Q5Q0T9 (Rat) > 97% Oligonucleotide synthesis is monitored base by base throug appropriate coupling efficiency. The oligo is subsequently pr phase extraction. The annealed RNA duplex is further analy spectrometry to verify the exact composition of the duplex. the previous lot by mass spectrometry to ensure maximum We offers pre-designed sets of 3 different target-specific siR METRN gene. Each vial contains 5 nmol of lyophilized siRNA transfected individually or pooled together to achieve knock gene, which is most commonly assessed by qPCR or western METRN siRNA (Rat) - A 5 nmol x 1			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

METRN siRNA (Rat) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com