

Product Data Sheet

TAAR7G siRNA (Rat)

Catalog #	Source	Reactivity	Applications		
CRR4336	Synthetic	R	RNAi		
Description	Description siRNA to inhibit TAAR7G expression using RNA interference				
Specificity	TAAR7	TAAR7G siRNA (Rat) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knock	down gene expression	on.		
Form	Lyoph	ilized powder			
Gene Symbol	TAAR7	TAAR7G			
Alternative N	ames TA9; T	TA9; TAR9; TRAR9; Trace amine-associated receptor 7g; TaR-7g; Trace amine			
	recept	tor 7g; Trace amine re	eceptor 9; TaR-9		
Entrez Gene	29413	1 (Rat)			
SwissProt	Q923)	Q923Y1 (Rat)			
Purity	> 97%	> 97%			
Quality Contr	ol Oligor	nucleotide synthesis i	ide synthesis is monitored base by base through trityl analysis to ensure		
	appro	priate coupling efficie	ency. The oligo is subsequently pur	ified by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spectr	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the pr	the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.			
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of rat			
	TAAR7	TAAR7G gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can be			
	transf	transfected individually or pooled together to achieve knockdown of the target			
	gene,	gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	TAAR	R7G siRNA (Rat) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

TAAR7G siRNA (Rat) - B	5 nmol x 1	5 nmol x 2
TAAR7G siRNA (Rat) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 μl
		10 nM	1 µl	5 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com